

INDICATIONS: For the prevention and treatment of anemia due to iron deficiency in baby pigs.

DOSAGE AND ADMINISTRATION: For intramuscular injection only.

Prevention: 1 mL (100 mg iron) at 2 to 4 days of age.

Treatment: 1 mL (100 mg iron).

Treatment may be repeated in 10 days.

NOTICE: Organic iron preparations injected intramuscularly into pigs beyond 4 weeks of age may cause staining of muscle tissue.

See accompanying leaflet for additional information.

LOT NO.:

EXP. DATE:

NDC 57319-494-05

Iron Hydrogenated Dextran Injection

Hematinic

**For Prevention and Treatment of
Iron Deficiency Anemia in Baby Pigs**

FOR ANIMAL USE ONLY

KEEP OUT OF REACH OF CHILDREN

**Net Contents:
100 mL**


PHOENIX™

Manufactured for:
Clipper Distributing Company, LLC
St. Joseph, MO 64507

Each mL of sterile solution contains 100 mg of elemental iron stabilized with a low molecular weight hydrogenated dextran and 0.5% phenol as a preservative.

Store at controlled room temperature between 15° and 30°C (59°–86°F).

Protect from freezing.

TAKE TIME



OBSERVE LABEL
DIRECTIONS

I-0900-04

Rev. 02-11

Trademarks are property
of Clipper Distributing
Company, LLC

Manufactured by
Sparhawk Laboratories, Inc.
Lenexa, KS 66215 USA

